

# ORGANOMETALLICS

Volume 17, Number 17, August 17, 1998

© Copyright 1998  
American Chemical Society

## Communications

### The Seven-Component Assembly of the Bowl-Shaped Cages $\text{Cp}^*\text{Rh}_7(\text{CN})_{12}^{2+}$ and $\text{Cp}^*\text{Rh}_3\text{Ir}_4(\text{CN})_{12}^{2+}$

Stephen M. Contakes, Kevin K. Klausmeyer, Richard M. Milberg,  
Scott R. Wilson, and Thomas B. Rauchfuss\*

School of Chemical Sciences, University of Illinois, Urbana, Illinois 61801


Received May 13, 1998

**Summary:** A novel bowl-shaped cyanometalate cage  $[\text{Cp}^*\text{Rh}_7(\text{CN})_{12}]^{2+}$  (**3**) forms via the reaction between MeCN solutions of  $\text{Et}_4\text{N}[\text{Cp}^*\text{Rh}(\text{CN})_3]$  and  $[\text{Cp}^*\text{Rh}(\text{NCMe})_3](\text{PF}_6)_2$ ; this seven-component condensation proceeds in 89% yield.  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopic measurements as well as a single-crystal X-ray diffraction studies show that **3** features a  $C_{3v}$   $\text{Rh}_7(\text{CN})_9$  cage, which can be described as a fragment of a  $\text{Rh}_8(\text{CN})_{12}$  box.

The prevailing interest in supramolecular coordination complexes,<sup>1,2</sup> together with recent advances in cyanometalates,<sup>3,4</sup> makes the study of cyanometalate cages topical and opportune. Most work in this area has focused on nonmolecular solids. Our recent synthesis<sup>5</sup> of the "molecular box",  $\text{Cp}_4\text{Cp}^*\text{Co}_4\text{Rh}_4(\text{CN})_{12}^{4+}$ , however, establishes the feasibility of constructing

molecular cages based on M–C–N–M linkages.<sup>6</sup> We now report results on a simple homometallic condensation process leading efficiently to organometallic species with bowl-shaped architectures.<sup>7,8</sup>

The key experiment involves the condensation of  $\text{Et}_4\text{N}[\text{Cp}^*\text{Rh}(\text{CN})_3]$  (**1**) and  $[\text{Cp}^*\text{Rh}(\text{NCMe})_3](\text{PF}_6)_2$  (**2**)<sup>9</sup> according to the ratios shown in eq 1.<sup>10</sup>



This transformation is conveniently monitored by  $^1\text{H}$  NMR spectroscopy, which, as independent studies show,<sup>11</sup> easily resolves  $\text{Cp}^*$  signals for  $\text{Rh}(\text{CN})_3$  ( $\sim\delta$  2.0) versus

(1) Lehn, J.-M. *Supramolecular Chemistry*; VCH: Weinheim, 1995.  
(2) Representative recent papers: Stang, P. J.; Olenyuk, B. *Acc. Chem. Res.* **1997**, *30*, 502. Fujita, M.; Ogura, K. *Coord. Chem. Rev.* **1996**, *148*, 249.

(3) Fehlhammer, W. P.; Fritz, M. *Chem. Rev.* **1993**, *93*, 1243. Dunbar, K. R.; Heitz, R. A. *Prog. Inorg. Chem.* **1997**, *45*, 283. Iwamoto, T.; Nishikiori, S.; Kitazawa, T.; Yuge, H. *J. Chem. Soc., Dalton Trans.* **1997**, 4197. Anwender, R. *Angew. Chem., Int. Ed.* **1998**, *37*, 599.

(4) Lu, J.; Paliwala, T.; Lim, S. C.; Yu, C.; Niu, T.; Jacobson, A. J. *Inorg. Chem.* **1997**, *36*, 923. Slone, R. V.; Yoon D. I.; Calhoun, R. M.; Hupp, J. T. *J. Am. Chem. Soc.* **1995**, *117*, 11813. Ramprasad, D.; Pez, G. P.; Toby, B. H.; Merkley, T. J.; Pearlstein, R. M. *J. Am. Chem. Soc.* **1995**, *117*, 10694. Entley, W. R.; Girolami, G. S. *Science* **1995**, *268*, 397. Brousseau, L. C.; Williams, D.; Kouvetakis, J.; O'Keeffe, M. *J. Am. Chem. Soc.* **1997**, *119*, 6292. Hoskins, B. F.; Robson, R. *J. Am. Chem. Soc.* **1990**, *112*, 1546.

(5) Klausmeyer, K. K.; Rauchfuss, T. B.; Wilson, S. R. *Angew. Chem., Int. Ed. Engl.* **1998**, *37*, 1705.

(6) For related cages based on triazacyclononane capping ligands in place of cyclopentadienyls: Heinrich, J. L.; Berseth, P. A.; Long, J. R. *Chem. Commun.* **1998**, 1231.


(7) Supramolecular organometallic chemistry based on  $\text{Cp}^*\text{Rh}^{2+}$ : Chen, H.; Ogo S.; Fish R. H. *J. Am. Chem. Soc.* **1996**, *118*, 4993, and references therein.

(8) Molecular bowls based on coordination compounds: Day, V. W.; Klemperer, W. G.; Yaghi, O. *J. Am. Chem. Soc.* **1989**, *111*, 5959. Holman, K. T.; Halihan, M. M.; Jurisson, S. S.; Atwood, J. L.; Burkhalter, R. S.; Mitchell, A. R.; Steed, J. W. *J. Am. Chem. Soc.* **1996**, *118*, 9567, and references therein.

(9) White, C.; Yates, A.; Maitlis, P. M. *Inorg. Synth.* **1992**, *29*, 228.

(10) A solution of 186 mg (0.414 mmol)  $\text{Et}_4\text{N}[\text{Cp}^*\text{Rh}(\text{CN})_3]$  in 10 mL of MeCN was added to a solution of 200 mg (0.307 mmol) of  $[\text{Cp}^*\text{Rh}(\text{MeCN})_3](\text{PF}_6)_2$  in 10 mL of MeCN. After 24 h, the reaction solution was concentrated to ca. 5 mL and diluted with 50 mL of  $\text{Et}_2\text{O}$  to give 240 mg of the product **3** (89%).



(11) Contakes, S. M.; Klausmeyer, K. K.; Rauchfuss, T. B. Unpublished results.



**Figure 1.** Structure of  $\text{Cp}^*_7\text{Rh}_7(\text{CN})_{12}^{2+}$  from two perspectives with thermal ellipsoids drawn at the 35% level. In the top perspective, the cation is viewed down the idealized 3-fold symmetry axis (molecules of solvation are not shown). In the bottom view, the  $\text{Rh}_7(\text{CN})_{12}$  core is shown together with the intracage solvent molecule (in this view, we included three  $\text{Cp}^*$  groups to highlight the degree to which the unique MeCN unit is inserted into the cage). Selected angles:  $\text{C}(1)-\text{Rh}(1)-\text{C}(1')$ ,  $94.0(4)^\circ$ ;  $\text{C}(1)-\text{Rh}(1)-\text{C}(2)$ ,  $88.7(2)^\circ$ ;  $\text{C}(4)-\text{Rh}(4)-\text{C}(6)$ ,  $88.9(3)^\circ$ ;  $\text{C}(3)-\text{Rh}(4)-\text{C}(4)$ ,  $87.0(2)^\circ$ ;  $\text{C}(3)-\text{Rh}(4)-\text{C}(6)$ ,  $87.8(3)^\circ$ ;  $\text{N}(1)-\text{Rh}(2)-\text{N}(3)$ ,  $88.3(2)^\circ$ ;  $\text{N}(1)-\text{Rh}(2)-\text{N}(5)$ ,  $87.1(2)^\circ$ ;  $\text{N}(3)-\text{Rh}(2)-\text{N}(5)$ ,  $90.6(2)^\circ$ ;  $\text{N}(2)-\text{Rh}(3)-\text{N}(4)$ ,  $88.9(2)^\circ$ .  $\text{Rh}-\text{N}-\text{C}$  angles range from  $169(2)^\circ$  for  $\text{C}(3)-\text{N}(3)-\text{Rh}(2)$  to  $178.7(7)^\circ$  for  $\text{C}(2)-\text{N}(2)-\text{Rh}(3)$ .  $\text{Rh}-\text{C}-\text{N}$  angles range from  $173.0(3)^\circ$  to  $178.5(5)^\circ$ .  $\text{Rh}-\text{N}$  distances range from  $2.065(5)$  to  $2.109(7)$  Å, while  $\text{Rh}-\text{C}(\text{N})$  distances range from  $1.985(7)$  to  $2.033(8)$  Å.

$\text{Rh}(\text{NC})_3$  ( $\sim\delta$  1.7) centers. Mixing MeCN solutions of **1** and **2** (4:3 ratio) results in the rapid (1 min) consumption of **2** and the formation of a complex mixture. After 1 day,  $[\text{Cp}^*_7\text{Rh}_7(\text{CN})_{12}](\text{PF}_6)_2$  (**3**) is the dominant species (89% isolated yield). Consistent with its crystallographically verified structure (see below), **3** has three  $^1\text{H}$  NMR signals with relative intensities of 1:3:3, the first two of which fall in the  $\text{Cp}^*\text{Rh}(\text{CN})_3$  region, while the other signal is in the  $\text{Cp}^*\text{Rh}(\text{NC})_3$  region.

The rate of assembly of **3** is comparable to that for the box  $\text{Cp}_4\text{Cp}^*_4\text{Co}_4\text{Rh}_4(\text{CN})_{12}^{4+}$  from  $[\text{K}(\text{18-crown-6})][\text{CpCo}(\text{CN})_3]$  and  $[\text{Cp}^*\text{Rh}(\text{NCMe})_3](\text{PF}_6)_2$ .<sup>11</sup> It is striking, however, that the all- $\text{Cp}^*$  box is never observed. Instead, equimolar quantities of **1** and **2** react to produce **3**, leaving some unreacted **2**. We suggest that the



**Figure 2.** 125 MHz  $^{13}\text{C}$  NMR spectrum of a MeCN solution of **3** (33%  $^{13}\text{C}$ -enriched in CN region).

formation of  $\text{Cp}^*_8\text{Rh}_8(\text{CN})_{12}^{4+}$  is prevented by repulsive  $\text{Cp}^*-\text{Cp}^*$  interactions.<sup>12</sup> Solvent also strongly affects the assembly process: in  $\text{CH}_2\text{Cl}_2$  solution, compounds **1** and **2** (4:3 ratio) react, but **3** is not formed until the solution is diluted with an equal volume of MeCN.

The formulation of **3** as  $[\text{Cp}^*_7\text{Rh}_7(\text{CN})_{12}](\text{PF}_6)_2$ <sup>13</sup> was confirmed by electrospray mass spectrometry on MeCN solutions, which showed peaks for the dicationic cage at  $m/z = 2128$  ( $z = 1$ ) and  $991$  ( $z = 2$ ). The solid-state structure of **3** was deduced by X-ray crystallographic analysis of crystals grown from MeCN/ $\text{Et}_2\text{O}$  (Figure 1). The dicationic cage has idealized  $\text{C}_{3v}$  symmetry with three types of  $\text{Cp}^*\text{Rh}$  centers.<sup>14</sup> Nine of the 12 CN ligands bridge between Rh atoms; the remaining three CN groups are terminal. The  $\text{C}-\text{Rh}-\text{C}$  and  $\text{N}-\text{Rh}-\text{N}$  angles range from  $87.1^\circ$  to  $94^\circ$ , i.e., **3** is well described as three conjoined  $\text{Rh}_4(\mu\text{-CN})_4$  squares. Thus, the  $\text{Rh}_7(\text{CN})_9$  core is a fragment of a  $\text{M}_8(\text{CN})_{12}$  box, which cannot form from **3** because three  $\text{CN}_{\text{term}}$  groups are oriented away from the cage interior. This change in stereochemistry, versus that of the cubic mixed  $\text{Cp}-\text{Cp}^*$  cage  $\text{Cp}_4\text{Cp}^*_4\text{Co}_4\text{Rh}_4(\text{CN})_{12}^{4+}$ , minimizes inter- $\text{Cp}^*$  steric repulsions which would prevail in a  $\text{Cp}^*_8\text{Rh}_8(\text{CN})_{12}^{4+}$

(12) Reaction of  $\text{Et}_3\text{N}[(\text{C}_5\text{Me}_4\text{H})\text{Rh}(\text{CN})_3]$  and  $[(\text{C}_5\text{Me}_4\text{H})\text{Rh}(\text{NCMe})_3](\text{PF}_6)_2$  in a 1:1 ratio also fails to give molecular boxes. However, use of a 4:3 ratio of reactants affords  $[(\text{C}_5\text{Me}_4\text{H})_7\text{Rh}_7(\text{CN})_{12}](\text{PF}_6)_2$  as the exclusive product.

(13)  $[\text{Cp}^*_7\text{Rh}_7(\text{CN})_{12}](\text{PF}_6)_2$ :  $^1\text{H}$  NMR ( $\text{CD}_3\text{CN}$  soln): 2.14 (s, 15H), 1.98 (s, 45H), and 1.70 (s, 45H).  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CD}_3\text{CN}$  soln, see Figure 2). ESI MS:  $m/z$  991.2 ( $\text{M}^{2+}$ ) 2128 ( $\text{M}^{2+} \text{PF}_6^-$ ) (mass spect. sample was prepared with 33%  $^{13}\text{C}$ ). IR (KBr,  $\text{cm}^{-1}$ ): 2126 ( $\nu_{\text{CN}}$ ) and 2169 ( $\nu_{\mu\text{-CN}}$ ). Anal. CHN.

box. The solid-state structure of **3** features a MeCN molecule inserted into the cage.<sup>15</sup> The inclusion capability of the cage is thus demonstrated, although the scope and energetics of this process will require further study.

As the connectivity of the CN ligands is highly relevant to the assembly process, we sought confirmatory evidence for the proposed structure beyond that afforded by the crystallographic analysis. To this end, we conducted one- and two-dimensional <sup>13</sup>C NMR measurements on samples of **3** prepared from 33% <sup>13</sup>C-enriched cyanide.<sup>16</sup> Observed were three sets (6:3:3) of CN signals, all of which showed strong coupling (51–56 Hz) to <sup>103</sup>Rh as well as additional longer range <sup>2</sup>J(<sup>13</sup>C, <sup>103</sup>Rh) and <sup>2</sup>J(<sup>13</sup>C, <sup>103</sup>Rh) fine structure (Figure 2). The three terminal CN groups absorb at  $\delta$  125, while the two types of  $\mu$ -CN groups, which can be assigned on the basis of their intensities, absorb in the region  $\delta$

(14) C<sub>90</sub>H<sub>123</sub>F<sub>12</sub>N<sub>16</sub>O<sub>3</sub>P<sub>2</sub>Rh<sub>7</sub>; fw = 2487.35;  $T = 175$  K;  $\lambda = 0.710$  73 Å; crystal size = 0.15 × 0.08 × 0.08 mm; Siemens CCD detector; orthorhombic (*Pnma*);  $a = 24.7243(6)$  Å,  $b = 16.8342(5)$  Å,  $c = 27.1790(7)$  Å;  $\alpha = \beta = \gamma = 90^\circ$ ;  $V = 11\,312.3(5)$  Å<sup>3</sup>;  $Z = 4$ ;  $\rho = 1.460$  Mg/m<sup>3</sup>; 7232 reflns ( $2 \leq \theta \leq 22$ ); rflns = 44 725 [ $R_{\text{int}} = 0.1883$ ] (indep rflns = 7232 [2169 obs,  $I > 2\sigma(I)$ ]); direct methods; data/restraints/parameters = 7227/457/578; GOF on  $F^2 = 1.100$ ; R1 (obs data) = 0.0871, wR2 = 0.1917; R1 (all data) = 0.1597, wR2 = 0.2393; largest diff peak and hole = 1.030 and  $-0.910$  e·Å<sup>-3</sup>. The deposition number with the Cambridge Crystallographic Data Centre is 101195.

(15) In the solid state, the cages in **3** are solvated by two molecules of H<sub>2</sub>O per formula unit; these water molecules are situated near (3.05 Å) two CN<sub>term</sub> ligands. In addition to the MeCN that is inside the cage, two other MeCN molecules/formula unit are positioned near two of the three Rh<sub>4</sub>(CN)<sub>4</sub> faces (in Cp<sub>4</sub>Cp\*<sub>4</sub>Co<sub>4</sub>Rh<sub>4</sub>(CN)<sub>12</sub><sup>4+</sup> each of the six faces is solvated by MeCN. See ref 5).

(16) <sup>125</sup> MHz <sup>13</sup>C NMR in CN region ( $J_s$  in Hz): 139.3 ( $J_{\text{C-Rh}} = 54$ , <sup>2</sup> $J_{\text{C-Rh}} = 3.6$ ), 136.5 ( $J_{\text{C-Rh}} = 56$ , <sup>2</sup> $J_{\text{C-Rh}} \sim 2$ ), 124.1 ( $J_{\text{C-Rh}} = 50$ , <sup>2</sup> $J_{\text{C-C}} = 10$ ). These assignments are consistent with the results of a <sup>13</sup>C COSY experiment.

136–140. These results prove that the cage assembles without cleavage of Rh–C bonds. This proposal is further supported by the synthesis of heterometallic analogues of **3**: the reaction of (Et<sub>4</sub>N)[Cp\*Ir(CN)<sub>3</sub>] (**4**) with **2** (4:3 ratio) afforded [Cp\*<sub>7</sub>Ir<sub>4</sub>Rh<sub>3</sub>(CN)<sub>12</sub>](PF<sub>6</sub>)<sub>2</sub> (**5**) in 69% isolated yield.<sup>17</sup>

In summary, this paper demonstrates the considerable potential of organometallic cyanometalates as precursors to novel cage structures. The formation of the new M<sub>7</sub>(CN)<sub>9</sub> cages results from a balance between inter-Cp\* steric repulsions and attractive M···N interactions.

**Acknowledgment.** This research is supported by the Department of Energy. S.M.C. thanks NSF for a Graduate Fellowship.

**Supporting Information Available:** Synthetic and characterization details for **1**, **3**, **4**, and **5**; details of the crystal structure analysis of **3** including tables of positions, displacement parameters, and bond angles and distances (22 pages). Ordering and Internet access information is given on any current masthead page.

OM9803772

(17) A solution of 107 mg (0.20 mmol) of Et<sub>4</sub>N[(Cp\*)Ir(CN)<sub>3</sub>] in 10 mL of MeCN was added to a solution of 98 mg (0.15 mmol) of [(Cp\*)Rh(MeCN)<sub>3</sub>](PF<sub>6</sub>)<sub>2</sub> in 10 mL of MeCN. After 24 h, the reaction solution was concentrated to ca. 5 mL and diluted with 50 mL of Et<sub>2</sub>O to precipitate the product **5** as a yellow powder. Yield: 206 mg (69%). {[Cp\*Ir(CN)<sub>3</sub>]<sub>4</sub>[Cp\*Rh<sub>3</sub>](PF<sub>6</sub>)<sub>2</sub>}. <sup>1</sup>H NMR (500 MHz, MeCN):  $\delta$  1.68 (s, 45H); 2.07 (s, 45H); 2.12 (s, 15H). <sup>13</sup>C NMR (125 MHz, MeCN):  $\delta$  9.92, 9.97, 10.40, 15.47, 66.14, 98.00, 98.20, 98.72, 99.23, 109.31, 112.86. ESI MS:  $m/z$  2480.9 (M – PF<sub>6</sub>) 1168.3 (M<sup>2+</sup>). IR (KBr, cm<sup>-1</sup>): 2247 ( $\nu_{\mu\text{-CN}}$ ) and 2170 ( $\nu_{\text{CN}}$ ). Anal. CHN.